

Order α_s^2 Contributions to the Fragmentation Functions in e^+e^- -Annihilation

W.L. van Neerven ¹

*Instituut-Lorentz,
University of Leiden,
PO Box 9506, 2300 RA Leiden,
The Netherlands.*

Abstract

We discuss the most important results which are obtained from a recent calculation of the order α_s^2 corrections to the coefficient functions contributing to the fragmentation functions $F_k(x, Q^2)$ ($k = T, L, A$) in e^+e^- -annihilation. From these fragmentation functions one can derive the corresponding integrated transverse (σ_T), longitudinal (σ_L) and asymmetric (σ_A) cross sections. The sum $\sigma_{tot} = \sigma_T + \sigma_L$ corrected up to order α_s^2 agrees with the well known result in the literature providing us with an independent check on our calculation. It turns out that the order α_s^2 corrections to the transverse and asymmetric quantities are small. However these corrections to $F_L(x, Q^2)$ and σ_L are large so that one gets a better agreement between the theoretical predictions and the data obtained from the LEP experiments.

¹talk presented at the Rencontres de Physique de La Valee d'Aoste, La Thuile, Italy, March 2-8, 1997

The measurement of the fragmentation functions in the process

$$e^+ + e^- \rightarrow \gamma, Z \rightarrow H + \text{“}X\text{”}, \quad (1)$$

provides us, in addition to other experiments like deep inelastic lepton-hadron scattering, with a new test of scaling violations as predicted by perturbative quantum chromodynamics (QCD). Here “ X ” denotes any inclusive final hadronic state and H represents either a specific charged outgoing hadron or a sum over all charged hadron species. This process has been studied over a wide range of energies of many different e^+e^- -colliders. The most recent data are coming from LEP (see [1],[2],[3]). The unpolarized differential cross section of the above process is given by

$$\frac{d^2\sigma^H}{dx d\cos\theta} = \frac{3}{8}(1 + \cos^2\theta)\frac{d\sigma_T^H}{dx} + \frac{3}{4}\sin^2\theta\frac{d\sigma_L^H}{dx} + \frac{3}{4}\cos\theta\frac{d\sigma_A^H}{dx}. \quad (2)$$

The Bjørken scaling variable x is defined by

$$x = \frac{2pq}{Q^2}, \quad q^2 = Q^2 > 0, \quad 0 < x \leq 1, \quad (3)$$

where p and q are the four-momenta of the produced particle H and the virtual vector boson (γ, Z) respectively. In the centre of mass (CM) frame of the electron-positron pair the variable x can be interpreted as a fraction of the beam energy carried away by the hadron H . The variable θ denotes the angle of emission of particle H with respect to the electron beam direction in the CM frame. The transverse, longitudinal and asymmetric cross sections in (2) are defined by σ_T^H , σ_L^H , and σ_A^H respectively. The latter only shows up if the intermediate vector boson is given by the Z -boson and is absent in purely electromagnetic annihilation.

Before the advent of LEP1 the CM energies were so low ($\sqrt{s} < M_Z$) that σ_A could not be measured and no effort was made to separate σ_T from σ_L so that only data for the sum of the above cross sections in eq. (2) were available. Recently after LEP1 came into operation ALEPH and OPAL obtained data for σ_T and σ_L separately and the latter collaboration even made a measurement of σ_A for the first time. The separation of σ_T and σ_L is important because

the latter cross section enables us to extract the strong coupling constant α_s and allows us to determine the gluon fragmentation density $D_g^H(x, \mu^2)$ with a much higher accuracy as could be done before. Furthermore the measurement of σ_A provides us with information on hadronization effects since the QCD corrections are very small. In the QCD improved parton model, which describes the production of the parton denoted by l ($l = q, \bar{q}, g$) and its subsequent fragmentation into hadron H , the cross sections σ_k^H ($k = T, L, A$) can be expressed as follows

$$\begin{aligned} \frac{d\sigma_k^H}{dx} = \int_x^1 \frac{dz}{z} \left[\sigma_{\text{tot}}^{(0)}(Q^2) \left\{ D_S^H \left(\frac{x}{z}, \mu^2 \right) \mathcal{C}_{k,q}^S(z, Q^2/\mu^2) + D_g^H \left(\frac{x}{z}, \mu^2 \right) \cdot \right. \right. \\ \left. \left. \cdot \mathcal{C}_{k,g}^S(z, Q^2/\mu^2) \right\} + \sum_{f=1}^{n_f} \sigma_f^{(0)}(Q^2) D_{NS,f}^H \left(\frac{x}{z}, \mu^2 \right) \mathcal{C}_{k,q}^{NS}(z, Q^2/\mu^2) \right], \quad (4) \end{aligned}$$

for $k = T, L$. In the case of the asymmetric cross section we have

$$\frac{d\sigma_A^H}{dx} = \int_x^1 \frac{dz}{z} \left[\sum_{f=1}^{n_f} A_f^{(0)}(Q^2) D_{A,f}^H \left(\frac{x}{z}, \mu^2 \right) \mathcal{C}_{A,q}^{NS}(z, Q^2/\mu^2) \right]. \quad (5)$$

In the formulae (4) and (5) we have introduced the following notations. The function $D_l^H(z, \mu^2)$ denotes the parton fragmentation density corresponding to the hadron of species H . Further we have defined the singlet (S) and non-singlet (NS, A) combinations of the quark fragmentation densities. They are given by

$$D_S^H(z, \mu^2) = \frac{1}{n_f} \sum_{q=1}^{n_f} \left(D_q^H(z, \mu^2) + D_{\bar{q}}^H(z, \mu^2) \right), \quad (6)$$

$$D_{NS,q}^H(z, \mu^2) = D_q^H(z, \mu^2) + D_{\bar{q}}^H(z, \mu^2) - D_S^H(z, \mu^2), \quad (7)$$

$$D_{A,q}^H(z, \mu^2) = D_q^H(z, \mu^2) - D_{\bar{q}}^H(z, \mu^2). \quad (8)$$

The index q stands for the quark species and n_f denotes the number of light flavours. Further the variable μ appearing in $D_l^H(z, \mu^2)$ and the coefficient

functions $\mathcal{C}_{k,l}(z, Q^2/\mu^2)$ ($k = T, L, A$; $l = q, g$) stands for the mass factorization scale which for convenience has been put equal to the renormalization scale. The pointlike cross section $\sigma_q^{(0)}$ and the asymmetry factor $A_q^{(0)}$ of the process $e^+ + e^- \rightarrow q + \bar{q}$ can be found in eq. (10) and eq. (15) of [4] respectively. The total cross section, summed over all flavours is given by $\sigma_{\text{tot}}^{(0)}(Q^2) = \sum_{q=1}^{n_f} \sigma_q^{(0)}(Q^2)$. The QCD corrections in eqs. (4), (5) are described by the coefficient functions $\mathcal{C}_{k,l}^r$ ($k = T, L, A$; $l = q, g$) which can be distinguished with respect to the flavour group $SU(n_f)$ into a singlet ($r = S$) and a non-singlet part ($r = NS$). The gluonic coefficient function only receives contributions from flavour singlet channel partonic subprocesses so that we can drop the superscript S on $\mathcal{C}_{k,g}$. However the quark coefficient functions get flavour singlet as well as flavour non-singlet contributions. Up to first order in the strong coupling constant α_s it turns out that $\mathcal{C}_{k,q}^{NS} = \mathcal{C}_{k,q}^S$. However in higher order both quantities start to deviate from each other. Hence we define the purely singlet coefficient function $\mathcal{C}_{k,q}^{PS}$ via

$$\mathcal{C}_{k,q}^S = \mathcal{C}_{k,q}^{NS} + \mathcal{C}_{k,q}^{PS}. \quad (9)$$

Like $\mathcal{C}_{k,g}$ the purely singlet coefficient function only receives contributions from the flavour singlet channel partonic subprocesses which in the former case show up for the first time in order α_s^2 .

Using charge conjugation invariance of the strong interactions one can show that $\mathcal{C}_{A,q}^{NS} = -\mathcal{C}_{A,\bar{q}}^{NS}$ and $\mathcal{C}_{A,q}^{PS} = \mathcal{C}_{A,g} = 0$. This implies that to σ_A^H in eq. (5) only non-singlet channel partonic subprocesses can contribute. Another important property of the coefficient function is that they do not depend on the probe γ or Z so that one can extract the overall pointlike cross section $\sigma_q^{(0)}$ or the asymmetry factor $A_q^{(0)}$. However this is only true if all quark masses are equal to zero and if one sums over all quark members in one family provided the latter appear in the inclusive state of the partonic subprocess (see [4]) From (2) we can derive the total hadronic cross section

$$\sigma_{\text{tot}}(Q^2) = \frac{1}{2} \sum_H \int_0^1 dx \int_{-1}^1 d \cos \theta \left(x \frac{d^2 \sigma^H}{dx d \cos \theta} \right) = \sigma_T(Q^2) + \sigma_L(Q^2), \quad (10)$$

with

$$\sigma_k(Q^2) = \frac{1}{2} \sum_H \int_0^1 dx x \frac{d\sigma_k^H}{dx}, \quad (k = T, L, A), \quad (11)$$

where one has summed over all types of outgoing hadrons H . From the momentum conservation sum rule given by

$$\sum_H \int_0^1 dx x D_l^H(x, \mu^2) = 1 \quad l = q, \bar{q}, g \quad (12)$$

and eqs. (4), (11) one can derive

$$\sigma_k(Q^2) = \sigma_{\text{tot}}^{(0)}(Q^2) \int_0^1 dx x \left[\mathcal{C}_{k,q}(x, Q^2/\mu^2)^S + \frac{1}{2} \mathcal{C}_{k,g}(x, Q^2/\mu^2) \right], \quad (13)$$

Finally we also define the transverse, longitudinal and asymmetric fragmentation functions $F_k^H(x, Q^2)^2$

$$F_k^H(x, Q^2) = \frac{1}{\sigma_{\text{tot}}^{(0)}(Q^2)} \frac{d\sigma_k^H}{dx}, \quad k = (T, L, A). \quad (14)$$

One observes that the above fragmentation functions are just the timelike analogues of the structure functions measured in for deep inelastic electron-proton scattering. The calculation of the coefficient functions corrected up to order α_s^2 proceeds in the same way as done for the Drell-Yan process in [5] and for deep inelastic lepton-hadron scattering in [6]. Denoting the intermediate vector bosons γ and Z by the symbol V we have the following parton subprocesses. In zeroth order we have the Born reaction

$$V \rightarrow "q" + \bar{q} \quad (15)$$

where " l " ($l = q, \bar{q}, g$) denotes the detected parton which subsequently fragments into the hadron of species H . In next-to-leading order (NLO) one has

²Notice that we make a distinction in nomenclature between the fragmentation densities D_q^H, D_g^H and the fragmentation functions F_k^H .

to include the one-loop virtual corrections to reaction (15) and the parton subprocesses

$$V \rightarrow "q" + \bar{q} + g \quad (16)$$

$$V \rightarrow "g" + q + \bar{q}. \quad (17)$$

After mass factorization of the collinear divergences which arise in the above processes one obtains the coefficient functions which are presented in [7],[8],[9]. The determination of the order α_s^2 contributions involves the computation of the two-loop corrections to (15) and the one-loop corrections to eqs. (16),(17). Furthermore one has to calculate the following subprocesses

$$V \rightarrow "q" + \bar{q} + g + g \quad (18)$$

$$V \rightarrow "g" + q + \bar{q} + g \quad (19)$$

$$V \rightarrow "q" + \bar{q} + q + \bar{q}. \quad (20)$$

In reaction (20)) the two anti-quarks, which are inclusive, can be identical as well as non identical. Notice that in the above reactions the detected quark can be replaced by the detected anti-quark so that in reaction (20) one can also distinguish between the final states containing identical quarks and non identical quarks. After mass factorization and renormalization for which we have chosen the $\overline{\text{MS}}$ -scheme one obtains the order α_s^2 contributions to the coefficient functions which are presented in [4],[10],[11]. The most important results of our calculations can be summarized as follows. From eq. (13) and the coefficient functions originating from the processes above we can obtain σ_L and σ_T corrected up to order α_s^2

$$\begin{aligned} \sigma_L(Q^2) = & \sigma_{\text{tot}}^{(0)}(Q^2) \left[\frac{\alpha_s(\mu^2)}{4\pi} C_F \left\{ 3 \right\} + \left(\frac{\alpha_s(\mu^2)}{4\pi} \right)^2 \left[C_F^2 \left\{ -\frac{15}{2} \right\} + C_A C_F \left\{ \right. \right. \right. \\ & \left. \left. \left. - 11 \ln \frac{Q^2}{\mu^2} - \frac{24}{5} \zeta(3) + \frac{2023}{30} \right\} + n_f C_F T_f \left\{ 4 \ln \frac{Q^2}{\mu^2} - \frac{74}{3} \right\} \right] \right], \end{aligned} \quad (21)$$

$$\begin{aligned}
\sigma_T(Q^2) = & \sigma_{\text{tot}}^{(0)}(Q^2) \left[1 + \left(\frac{\alpha_s(\mu^2)}{4\pi} \right)^2 \left[C_F^2 \left\{ 6 \right\} + C_A C_F \left\{ -\frac{196}{5} \zeta(3) - \frac{178}{30} \right\} \right. \right. \\
& \left. \left. + n_f C_F T_f \left\{ 16 \zeta(3) + \frac{8}{3} \right\} \right] \right]. \quad (22)
\end{aligned}$$

Addition of σ_L and σ_T yields the well known answer for σ_{tot} (10) which is in agreement with the literature [12]. Hence this quantity provides us with a check on our calculation of the longitudinal and transverse coefficient functions. Notice that in lowest order σ_{tot} only receives a contribution from the transverse cross section whereas the order α_s contribution can be only attributed to the longitudinal part. In order α_s^2 both σ_L and σ_T contribute to σ_{tot} .

Because of the high sensitivity of expression (21) to the value of α_s , the longitudinal cross section provides us with an excellent tool to measure the running coupling constant.

To illustrate the sensitivity of σ_L on α_s we have plotted in Fig. 1 in LO and NLO the ratio

$$R_L(Q^2) = \frac{\sigma_L(Q^2)}{\sigma_{\text{tot}}(Q^2)} \quad (23)$$

as a function of Q (CM-energy of the e^+e^- system). Our input parameters for the running coupling constant are $\Lambda_{LO}^{(5)} = 108 \text{ MeV}$ ($\alpha_s^{LO}(M_Z) = 0.122$) and $\Lambda_{\overline{\text{MS}}}^{(5)} = 227 \text{ MeV}$ ($\alpha_s^{NLO}(M_Z) = 0.118$). Fig. 1 reveals that the order α_s corrections are appreciable and they vary from 48% ($Q = 10 \text{ GeV}$) down to 28% ($Q = 200 \text{ GeV}$) with respect to the LO result. Furthermore one observes an improvement of the renormalization scale dependence while going from R_L^{LO} to R_L^{NLO} . Comparing with the experimental value $R_L = 0.057 \pm 0.005$, measured by OPAL [3], one observes a considerable improvement when the order α_s^2 contributions are included. However there is still a little discrepancy between R_L^{NLO} at $\mu = Q = M_Z$, and the data. This can either be removed by choosing a larger $\Lambda_{\overline{\text{MS}}}$ or by including the masses of the heavy quarks c and b in the calculation of the coefficient functions. Also a contribution of higher twist effects can maybe not neglected (see [9]).

We now want to investigate the effect of the order α_s^2 contributions to the

longitudinal and transverse fragmentation functions $F_L(x, Q^2)$ and $F_T(x, Q^2)$ eq. (14). Here we have summed over the following hadron species i.e. $H = \pi^\pm, K^\pm, p, \bar{p}$. Choosing the parametrization of the fragmentation densities in [13], with $\Lambda_{LO}^{(5)} = \Lambda_{\overline{\text{MS}}}^{(5)} = 190 \text{ MeV}$, we have plotted F_L^{LO} and F_L^{NLO} in Fig. 1. We observe that F_L^{LO} is below the data in particular in the small x -region. The agreement with the data becomes better when the order α_s corrections are included although at very small x F_L^{NLO} is still smaller than the values given by experiment. This figure reveals the importance of the higher order corrections to $F_L(x, Q^2)$ for the determination of the fragmentation densities. We have also shown results for F_T^{NLO} and F_T^{NNLO} in Fig. 3 using the same fragmentation density set. Both fragmentation functions agree with the data except that F_T^{NNLO} gets a little bit worse at very small x . Furthermore F_T^{NLO} and F_T^{NNLO} hardly differ from each other which means that the order α_s^2 corrections are small. One also notices that F_L constitutes the smallest part of the total fragmentation function $F = F_T + F_L$ which can be inferred from Figs. 2 and 3. This in particular holds at large x where $F_T \gg F_L$. Hence a fit of the fragmentation densities to the data of F_T is not sufficient to give a precise prediction for F_L and one has to include the data of the longitudinal part to provide us with a better set of fragmentation densities. This in particular holds for $D_g^H(z, \mu^2)$ in the small z -region for which the order α_s^2 contribution to F_L will be needed. The latter will certainly change the parametrization of the gluon fragmentation density given by ALEPH in [2] and OPAL in [3].

References

- [1] P. Abreu et al. (DELPHI), Phys. Lett. 311B (1993) 408.
- [2] D. Buskulic et al. (ALEPH), Phys. Lett. 357B (1995) 487.
- [3] R. Akers et al. (OPAL), Z. Phys. C68 (1995) 203.
- [4] P.J. Rijken and W.L. van Neerven, Nucl. Phys. 487B (1997) 233.

- [5] R. Hamberg, W.L. van Neerven, T. Matsuura, Nucl. Phys. B359 (1991) 343.
- [6] E.B. Zijlstra and W.L. van Neerven, Nucl. Phys. B383 (1992) 525.
- [7] P. G. Altarelli, R.K. Ellis, G. Martinelli, S.-Y. Pi, Nucl. Phys. B160 (1979) 301.
- [8] R. Baier and K. Fey, Z. Phys. C2 (1979) 339.
- [9] P. Nason and B.R. Webber, Nucl. Phys. B421 (1993) 473.
- [10] P.J. Rijken and W.L. van Neerven, Phys. Lett. B386 (1996) 422.
- [11] P.J. Rijken and W.L. van Neerven, Phys. Lett. 392B (1997) 207.
- [12] K.G. Chetyrkin, A.L. Kataev, F.V. Tkachov, Phys. Lett. 85B (1979) 277;
M. Dine and J. Sapirstein, Phys. Rev. Lett. 43 (1979) 668;
W. Celmaster and R.J. Gonsalves, Phys. Rev. Lett. 44 (1980) 560.
- [13] J. Binnewies, B.A. Kniehl, G. Kramer, Z. Phys. C65 (1995) 471.

Figure Captions

- Fig. 1** The ratio $R_L = \sigma_T/\sigma_{\text{tot}}$. Dotted lines: R_L^{LO} ; solid lines: R_L^{NLO} . Lower curve: $\mu = 2Q$; middle curve: $\mu = Q$; upper curve: $\mu = Q/2$. The data point at $Q = M_Z$ is from OPAL [3].
- Fig. 2** The longitudinal fragmentation function $F_L(x, Q^2)$ at $\mu = Q = M_Z$. Dotted line: F_L^{LO} ; solid line: F_L^{NLO} . The data are from ALEPH [2] and OPAL [3]. The fragmentation density set is from [13].
- Fig. 3** The transverse fragmentation function $F_T(x, Q^2)$ at $\mu = Q = M_Z$. Dotted line: F_T^{NLO} ; solid line: F_T^{NNLO} . The data are from ALEPH [2] and OPAL [3]. The fragmentation density set is from [13].




Figure 1:




Figure 2:




Figure 3: